

Public Health and High Traffic Roadways

California Air Resources Board Recommended Policy:

Sensitive land uses such as residences, schools, day care centers, playgrounds, and medical facilities should not be sited within 500 feet of:

- A freeway
- Urban roads with 100,000 or more vehicles/day
- Rural roads with 50,000 or more vehicles/day

(Ref. *"Air Quality and Land Use: A Community Health Perspective."* California Air Resources Board. April 2005)

Reason for the Policy:

Many studies show that living in proximity to freeways and other high traffic roadways leads to adverse health effects beyond those associated with regional air pollution. A number of studies that focused on children have found slower lung development and significant increases in the incidence of lung disease, such as asthma, bronchitis, and decreased lung function, in children who live or attend school near heavily travelled roadways. In addition to children, seniors, and people with heart and lung conditions are considered particularly sensitive to effects of air pollution. Residence in high-traffic areas has been shown to increase the risk of mortality within a cohort of male veterans.

Health Studies:

The results of health studies suggests that it is important to avoid exposing children and other sensitive populations to the elevated air pollution levels near freeways and other high traffic roads. While particulate pollution is suspected as contributing the most to the adverse health effects, studies have not yet determined which specific pollutants and sources (cf. diesel particulate, re-entrained roadway dust particulate, NO₂ vehicle exhaust, diesel trucks vs. gasoline cars, &c.) are responsible. Additional studies are underway. While significant adverse health effects were observed in children who lived within 1,500 feet of a freeway (Gauderman, 2007), the studies indicate a substantial benefit to a 500 foot separation (McConnell, 2006).

Key Findings:

- Reduced lung function in children is associated with traffic density within 1,000 feet and the strongest association is within 300 feet of the roadway. (Brunekreef, 1997)
- Children living within 550 feet of heavy traffic have more medical visits than children who live further away from traffic. (English, 1999)
- Increased asthma hospitalizations are associated with living within 650 feet of heavy traffic. (Lin, 2000)
- Asthma symptoms increase with proximity to roadways and the risk is greatest within 300 feet. (Venn, 2001)
- Asthma and bronchitis symptoms in children are associated with proximity to high traffic in a community with good overall regional air quality. (Kim, 2004)
- Children living within 150 – 200 meters (~450 feet – 600 feet) of heavy traffic have higher rates of asthma than children living further away from traffic. (McConnell, 2006)
- Children living within 500 meters (~1,500 feet) of heavy traffic have significantly slower lung development than children living further away from traffic. (Gauderman, 2007)
- Survival of members of the Washington University-EPRI Veterans Cohort is strongly and robustly associated with county-average levels of traffic related air pollution and mortality relationships are stronger in the counties with higher levels of traffic density. (Lipfert et al, 2009)

Applicability to Santa Barbara County:

The studies covered children in a variety of urban environments living in proximity to roadways covering a wide spectrum of traffic volumes. The adverse health effects were measured at traffic volumes as low as 41,000 vehicles per day (English) and between 80,000 and 150,000 vehicles per day (Brunekreef). Highway 101, through Santa Barbara County, experiences traffic volumes within the range where health effects have been observed. Also, some parts of Highway 101 see over 7000 diesel trucks per day

(SBCAG). Furthermore, running parallel to Highway 101 through the southern portion of Santa Barbara County is a rail corridor that contributes significantly to the pollution levels near the highway (cf., rail contributes an additional 10% or 0.07 tons per day to mobile source generated PM10 emissions in Santa Barbara County).

2012 Average Daily Traffic (ADT) Volumes for Highway 101 (Caltrans):

US 101 at Storke = 63,600 ADT (Back)

US 101 at Highway 150 = 64,000 ADT

US 101 at Las Positas/Route 225 = 130,000 ADT

US 101 at Highway 166, Santa Maria = 57,400 ADT

Conclusion:

In order to protect the public health, especially the health of children, from the adverse effects of air pollutants generated by traffic on Highway 101, land use policies should prohibit the construction of new residences, schools, day care centers, playgrounds, and medical facilities within 500 feet of Highway 101. No other roadways in Santa Barbara County currently have estimated traffic volumes at the magnitude for which the proximity studies have identified adverse health effects.

References:

- “Air Quality and Land Use: A Community Health Perspective.”* California Air Resources Board. (April 2005).
- W. James Gauderman, et al. *“Effect of exposure to traffic on lung development from 10 to 18 years of age: A cohort study.”* The Lancet. Volume 369, Issue 9561. 17 February 2007 – 23 February 2007: Pages 571-577.
- Rob McConnell, et al. *“Traffic, Susceptibility, and Childhood Asthma.”* Environmental Health Perspectives. Volume 114, Number 5, May 2006.
- John M. Peters, M.D., Sc.D. *“Epidemiologic Investigation to Identify Chronic Effects of Ambient Air Pollutants in Southern California (USC Children’s Health Study).”* California Air Resources Board (May 2004).
- Brunekreef, B. et al. *“Air pollution from truck traffic and lung function in children living near motorways.”* Epidemiology. 1997; 8:298-303.
- Lin, S. et al. *“Childhood asthma hospitalization and residential exposure to state route traffic.”* Environ Res. 2002;88:73-81.
- Venn. et al. *“Living near a main road and the risk of wheezing illness in children.”* American Journal of Respiratory and Critical Care Medicine. 2001; Vol.164, pp. 2177-2180.
- Kim, J. et al. *“Traffic-related air pollution and respiratory health: East Bay Children’s Respiratory Health Study.”* American Journal of Respiratory and Critical Care Medicine 2004; Vol. 170. pp. 520-526
- Zhu, Y et al. *“Study of Ultra-Fine Particles Near A Major Highway With Heavy-Duty Diesel Traffic.”* Atmospheric Environment. 2002 ; 36:4323-4335.
- Knape, M. *“Traffic related air pollution in city districts near motorways.”* The Science of the Total Environment. 1999; 235:339-341.
- Roseville Rail Yard Study.* California Air Resources Board (October 2004).
- ARB Diesel Risk Reduction Plan.* California Air Resources Board (2000).
- Delfino RJ *“Epidemiologic Evidence for Asthma and Exposure to Air Toxics: Linkages Between Occupational, Indoor, and Community Air Pollution Research.”* Environmental Health Perspectives. (2002) 110 (supplement 4): 573-589.
- English P., Neutra R., Scalf R. Sullivan M. Waller L. Zhu L. *“Examining Associations between Childhood Asthma and Traffic Flow Using a Geographic Information System.”* (1999) Environmental Health Perspectives 107(9): 761-767.
- “2007 Clean Air Plan.”* Santa Barbara County Air Pollution Control District (August 2007).
- “2007 Travel Trends Report for Santa Barbara County.”* Santa Barbara County Association of Governments. (December 2007)
- Lipfert, F. et al. *“Air Pollution and Survival within the Washington University-EPRI Veterans Cohort: Risks Based on Modeled Estimates of Ambient Levels of Hazardous and Criteria Air Pollutants.”* Journal of the Air & Waste Management Association. Vol. 59 April 2009, pp. 473-487.